

# THE KURATOWSKI COVERING CONJECTURE FOR GRAPHS OF ORDER $< 10$ FOR THE NONORIENTABLE SURFACES OF GENUS 3 AND 4

SUHKJIN HUR

ABSTRACT. Kuratowski proved that a finite graph embeds in the plane if it does not contain a subdivision of either  $K_5$  or  $K_{3,3}$ , called Kuratowski subgraphs. A conjectured generalization of this result to all nonorientable surfaces says that a finite minimal forbidden subgraph for the nonorientable surface of genus  $\tilde{g}$  can be written as the union of  $\tilde{g} + 1$  Kuratowski subgraphs such that the union of each pair of these fails to embed in the projective plane, the union of each triple of these fails to embed in the Klein bottle if  $\tilde{g} \geq 2$ , and the union of each triple of these fails to embed in the torus if  $\tilde{g} \geq 3$ . We show that this conjecture is true for all minimal forbidden subgraphs of order  $< 10$  for the nonorientable surfaces of genus 3 and 4.

## 1. INTRODUCTION

We use the same terminology as in [12] unless otherwise specified.  $G_1 \vee G_2$  denotes the graph obtained by identifying one vertex of  $G_1$  and one vertex of  $G_2$ .

Kuratowski [11] showed that minimal forbidden subgraphs for the plane are  $K_5$  and  $K_{3,3}$ . Given a graph  $G$ , any subgraph of  $G$  that is a subdivision of  $K_5$  or  $K_{3,3}$  is called a *Kuratowski subgraph* of  $G$ .

Then one might ask if Kuratowski's result can be extended to higher genus surfaces in terms of Kuratowski subgraphs. Glover has conjectured that if a finite graph  $G$  is a minimal forbidden subgraph for the nonorientable surface  $\mathbb{N}_{\tilde{g}}$ , then  $G$  can be written as the union of  $\tilde{g} + 1$  Kuratowski subgraphs such that the union of each pair of these fails to embed in the projective plane, the union of each triple of these fails to embed in the Klein bottle if  $\tilde{g} \geq 2$ , and the union of each triple of these fails to embed in the torus if  $\tilde{g} \geq 3$ . It should be noted that  $G$  is the union of  $\tilde{g} + 1$  Kuratowski subgraphs, i.e., every edge in  $G$  is an edge in at least one of the Kuratowski subgraphs. The set of  $\tilde{g} + 1$  subgraphs described in the conjecture is called a *Kuratowski covering* and the conjecture is called the *Kuratowski covering conjecture*. In this paper, we prove the following restricted version of the above conjecture.

**Theorem 1.1.** *The Kuratowski covering conjecture is true for every graph of order  $< 10$  for  $\mathbb{N}_3$  and  $\mathbb{N}_4$ .*

We prove this theorem by providing a Kuratowski covering for every minimal forbidden subgraph of order  $< 10$  for  $\mathbb{N}_3$  and  $\mathbb{N}_4$ .

We use the complete lists of minimal forbidden subgraphs of order  $< 10$  for some surfaces. Archdeacon and Huneke [2] showed that there are finitely many minimal forbidden subgraphs for nonorientable surfaces, and Robertson and Seymour [15] independently showed that there are finitely many minimal forbidden subgraphs for

arbitrary surfaces. Kuratowski [11] showed that minimal forbidden subgraphs for the plane are  $K_5$  and  $K_{3,3}$ . A list of minimal forbidden subgraphs for the projective plane has been found by Glover, Huneke, and Wang [5] and Archdeacon [1] proved that this list is complete. However, for the higher genus surfaces, the complete lists of minimal forbidden subgraphs are not known. The complete list of 8-vertex minimal forbidden subgraphs for the Klein bottle has been found by Huneke, McQuillan, and Richter [7] and the complete list of 9-vertex minimal forbidden subgraphs for the Klein bottle have been found by Cashy [3] and Hur [10]. The complete list of 8-vertex minimal forbidden subgraphs for the torus has been found by Duke and Haggard [4], and the complete list of 9-vertex minimal forbidden subgraphs for the torus has been found by Hlavacek [6]. We note that Wendy Myrvold in the Department of Computer Science, University of Victoria independently found about 235,000 minimal forbidden subgraphs for the torus using a computer [13], [14]. In particular, the list of 9-vertex minimal forbidden subgraphs for the torus found by Hlavacek coincides with the list of 9-vertex minimal forbidden subgraphs for the torus found by Myrvold [13].

The remainder of this paper is organized as follows. In Section 2, we find Kuratowski coverings for all minimal forbidden subgraphs of order  $< 10$  for  $\mathbb{N}_3$  and  $\mathbb{N}_4$ , which prove Theorem 1.1.

*Remark 1.* Theorem 1.1 is part of the main result of author's thesis [10], [9] : Every minimal forbidden subgraph of order  $< 10$  satisfies the Kuratowski covering conjecture.

*Remark 2.* A strengthened form of the Kuratowski covering conjecture analogous to the complete Kuratowski theorem for the plane says that a finite graph  $G$  fails to embed in  $\mathbb{N}_{\tilde{g}}$  if and only if there are  $\tilde{g} + 1$  Kuratowski subgraphs in  $G$  satisfying the conditions of the Kuratowski covering conjecture.











## 2. KURATOWSKI COVERINGS FOR $\mathbb{N}_3$ AND $\mathbb{N}_4$

**2.1. Kuratowski coverings for  $\mathbb{N}_3$ .** The complete list of minimal forbidden subgraphs of order  $< 10$  for  $\mathbb{N}_3$  is given in [8]. Since the genus of  $\mathbb{N}_3$  is three, for each minimal forbidden subgraph  $G$  of order  $< 10$  for  $\mathbb{N}_3$ , we find four Kuratowski subgraphs  $G_1, G_2, G_3, G_4$  as a Kuratowski covering such that the union of every pair of these contains a subdivision of a minimal forbidden subgraph for the projective plane and the union of every triple of these contains a subdivision of a minimal forbidden subgraph for the torus and a subdivision of a minimal forbidden subgraph for the Klein bottle. We show these Kuratowski coverings in Figure 1, ..., 34. The names of minimal forbidden subgraphs for the projective are from [5]. The 8-vertex minimal forbidden subgraphs for the torus are  $K_8 - K_3$ ,  $K_8 - (K_{1,2} \cup 2K_2)$ , and  $K_8 - K_{2,3}$  [4], and the 8-vertex minimal forbidden subgraphs for the Klein bottle are  $K_8 - 4K_2$ ,  $K_8 - (K_3 \vee K_2)$ ,  $K_8 - 2K_3$ ,  $K_8 - 2K_{1,3}$ , and  $K_8 - (K_{1,4} \cup K_3)$ , which are from [7]. The names of 9-vertex minimal forbidden subgraphs for the torus are from [6] and there are 63 9-vertex minimal forbidden subgraphs  $\tilde{I}_{9,1}^2, \dots, \tilde{I}_{9,63}^2$  for the Klein bottle [3], [10].


**2.2. Kuratowski coverings for  $\mathbb{N}_4$ .** There are two minimal forbidden subgraphs  $K_9 - K_{1,2}$  and  $K_9 - 2K_2$  of order  $< 10$  for  $\mathbb{N}_4$  [8] and we need five Kuratowski subgraphs  $G_1, G_2, G_3, G_4, G_5$  as a Kuratowski covering which satisfies the conditions

of the Kuratowski covering conjecture. We show Kuratowski coverings of  $K_9 - K_{1,2}$  and  $K_9 - 2K_2$  in Figure 35, ..., 40.


$$G = K_8 -$$

G<sub>1</sub>G<sub>2</sub>G<sub>3</sub>G<sub>4</sub>G<sub>1</sub> ∪ G<sub>2</sub> contains B<sub>1</sub>G<sub>2</sub> ∪ G<sub>3</sub> contains D<sub>3</sub>G<sub>1</sub> ∪ G<sub>3</sub> contains D<sub>3</sub>G<sub>2</sub> ∪ G<sub>4</sub> contains B<sub>4</sub>G<sub>1</sub> ∪ G<sub>4</sub> contains D<sub>17</sub>G<sub>3</sub> ∪ G<sub>4</sub> contains D<sub>3</sub>FIGURE 1. Kuratowski covering of  $G = \tilde{I}_{8,1}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
 $K_8 - (K_{1,2} \cup 2K_2)$  and  $K_8 - 4K_2$



$G_1 \cup G_3 \cup G_4$  contains  
 $K_8 - (K_{1,2} \cup 2K_2)$  and  $K_8 - (K_3 \vee K_2)$



$G_1 \cup G_2 \cup G_4$  is isomorphic  
to  $G_1 \cup G_3 \cup G_4$



$G_1 \cup G_2 \cup G_3$  contains  
 $K_8 - K_{2,3}$  and  $K_8 - (K_3 \vee K_2)$




FIGURE 2. Kuratowski covering of  $G = \tilde{I}_{8,1}^3$  for  $\mathbb{N}_3$  (Continued)






FIGURE 3. Kuratowski covering of  $G = \tilde{I}_{9,1}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_3$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$





FIGURE 4. Kuratowski covering of  $G = \tilde{I}_{9,1}^3$  for  $\mathbb{N}_3$  (Continued)

$$G = K_9 - \begin{array}{c} \bullet & \bullet \\ \diagup \quad \diagdown \\ \bullet \end{array}$$




G<sub>1</sub>


G<sub>2</sub>

G<sub>3</sub>


G<sub>4</sub>




$G_1 \cup G_2$  contains  $D_3$




$G_2 \cup G_3$  contains  $D_{17}$




$G_1 \cup G_3$  contains  $C_7$



$G_2 \cup G_4$  contains a subdivision of  $D_{17}$




$G_1 \cup G_4$  contains a subdivision of  $D_3$




$G_3 \cup G_4$  contains  $E_3$

FIGURE 5. Kuratowski covering of  $G = \tilde{I}_{9,2}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
S5.6 and  $\tilde{I}_{9,48}^2$



$G_1 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_3$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$




FIGURE 6. Kuratowski covering of  $G = \tilde{I}_{9,2}^3$  for  $\mathbb{N}_3$  (Continued)






FIGURE 7. Kuratowski covering of  $G = \tilde{I}_{9,3}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
T6.8 and  $\tilde{I}_{9,23}^2$



$G_1 \cup G_3 \cup G_4$  contains  
T6.8 and  $\tilde{I}_{9,23}^2$



$G_1 \cup G_2 \cup G_4$  contains  
S5.6 and  $\tilde{I}_{9,39}^2$



$G_1 \cup G_2 \cup G_3$  contains  
S5.6 and  $\tilde{I}_{9,39}^2$















FIGURE 8. Kuratowski covering of  $G = \tilde{I}_{9,3}^3$  for  $\mathbb{N}_3$  (Continued)


$$G = K_9 -$$

G<sub>1</sub>G<sub>2</sub>G<sub>3</sub>G<sub>4</sub>G<sub>1</sub> ∪ G<sub>2</sub> contains B<sub>2</sub>G<sub>2</sub> ∪ G<sub>3</sub> contains a subdivision of D<sub>3</sub>G<sub>1</sub> ∪ G<sub>3</sub> contains D<sub>7</sub>G<sub>2</sub> ∪ G<sub>4</sub> contains D<sub>6</sub>G<sub>1</sub> ∪ G<sub>4</sub> contains a subdivision of D<sub>17</sub>G<sub>3</sub> ∪ G<sub>4</sub> contains a subdivision of D<sub>3</sub>FIGURE 9. Kuratowski covering of  $G = \tilde{I}_{9,4}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_4$  contains  
U6.6a and  $\tilde{I}_{9,42}^2$



$G_1 \cup G_2 \cup G_3$  contains  
U6.6a and  $\tilde{I}_{9,42}^2$







FIGURE 10. Kuratowski covering of  $G = \tilde{I}_{9,4}^3$  for  $N_3$  (Continued)



$G_1 \cup G_2$  contains a subdivision of  $B_1$      $G_2 \cup G_3$  contains a subdivision of  $D_3$




$G_1 \cup G_3$  contains a subdivision of  $D_3$      $G_2 \cup G_4$  contains a subdivision of  $B_4$




$G_1 \cup G_4$  contains a subdivision of  $D_{17}$      $G_3 \cup G_4$  contains  $D_3$

FIGURE 11. Kuratowski covering of  $G = \tilde{I}_{9,5}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
S5.6 and  $\tilde{I}_{9,37}^2$



$G_1 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_4$  contains  
V6.5 and  $\tilde{I}_{9,39}^2$



$G_1 \cup G_2 \cup G_3$  contains  
U6.6a and  $\tilde{I}_{9,41}^2$















FIGURE 12. Kuratowski covering of  $G = \tilde{I}_{9,5}^3$  for  $N_3$  (Continued)


$$G = K_9 - \begin{array}{c} 1 \\ 4 \\ 7 \\ \swarrow \quad \searrow \\ 3 \quad 6 \\ \swarrow \quad \searrow \\ 2 \quad 5 \\ \swarrow \quad \searrow \\ 8 \end{array}$$

G<sub>1</sub>G<sub>2</sub>G<sub>3</sub>G<sub>4</sub>G<sub>1</sub> ∪ G<sub>2</sub> contains B<sub>5</sub>G<sub>2</sub> ∪ G<sub>3</sub> contains a subdivision of D<sub>3</sub>G<sub>1</sub> ∪ G<sub>3</sub> contains D<sub>3</sub>G<sub>2</sub> ∪ G<sub>4</sub> contains D<sub>3</sub>G<sub>1</sub> ∪ G<sub>4</sub> contains a subdivision of D<sub>3</sub>G<sub>3</sub> ∪ G<sub>4</sub> contains B<sub>5</sub>FIGURE 13. Kuratowski covering of  $G = \tilde{I}_{9,6}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_3$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$















FIGURE 14. Kuratowski covering of  $G = \tilde{I}_{9,6}^3$  for  $N_3$  (Continued)


$$G = K_9 -$$

G<sub>1</sub>G<sub>2</sub>G<sub>3</sub>G<sub>4</sub>G<sub>1</sub> ∪ G<sub>2</sub> contains B<sub>2</sub>G<sub>2</sub> ∪ G<sub>3</sub> contains a subdivision of D<sub>3</sub>G<sub>1</sub> ∪ G<sub>3</sub> contains D<sub>8</sub>G<sub>2</sub> ∪ G<sub>4</sub> contains E<sub>21</sub>G<sub>1</sub> ∪ G<sub>4</sub> contains a subdivision of D<sub>17</sub>G<sub>3</sub> ∪ G<sub>4</sub> contains a subdivision of D<sub>3</sub>FIGURE 15. Kuratowski covering of  $G = \tilde{I}_{9,7}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_4$  contains  
U6.6a and  $\tilde{I}_{9,42}^2$



$G_1 \cup G_2 \cup G_3$  contains  
U6.6a and  $\tilde{I}_{9,42}^2$















FIGURE 16. Kuratowski covering of  $G = \tilde{I}_{9,7}^3$  for  $N_3$  (Continued)


$$G = K_9 - \begin{array}{c} 2 \\ 9 \\ 3 \\ \diagdown \\ \diagup \\ 7 \\ 6 \\ 8 \\ 5 \\ 4 \end{array}$$

G<sub>1</sub>G<sub>2</sub>G<sub>3</sub>G<sub>4</sub>G<sub>1</sub> ∪ G<sub>2</sub> contains C<sub>7</sub>G<sub>2</sub> ∪ G<sub>3</sub> contains D<sub>7</sub>G<sub>1</sub> ∪ G<sub>3</sub> contains a subdivision of D<sub>3</sub>G<sub>2</sub> ∪ G<sub>4</sub> contains E<sub>21</sub>G<sub>1</sub> ∪ G<sub>4</sub> contains E<sub>20</sub>G<sub>3</sub> ∪ G<sub>4</sub> contains a subdivision of D<sub>3</sub>FIGURE 17. Kuratowski covering of  $G = \tilde{I}_{9,8}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_3$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$







FIGURE 18. Kuratowski covering of  $G = \tilde{I}_{9,8}^3$  for  $N_3$  (Continued)

FIGURE 19. Kuratowski covering of  $G = \tilde{I}_{9,9}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
S5.6 and  $\tilde{I}_{9,50}^2$



$G_1 \cup G_3 \cup G_4$  contains  
S5.6 and  $\tilde{I}_{9,50}^2$



$G_1 \cup G_2 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_3$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$














FIGURE 20. Kuratowski covering of  $G = \tilde{I}_{9,9}^3$  for  $N_3$  (Continued)


$$G = K_9 - \begin{array}{c} 1 \\ \diagdown \\ 3 \\ \diagup \\ 2 \end{array} \quad \begin{array}{c} 5 \\ \diagup \\ 4 \\ \diagdown \\ 6 \end{array}$$

G<sub>1</sub>G<sub>2</sub>G<sub>3</sub>G<sub>4</sub>G<sub>1</sub>  $\cup$  G<sub>2</sub> contains C<sub>7</sub>G<sub>2</sub>  $\cup$  G<sub>3</sub> contains a subdivision of B<sub>1</sub>G<sub>1</sub>  $\cup$  G<sub>3</sub> contains C<sub>7</sub>G<sub>2</sub>  $\cup$  G<sub>4</sub> contains C<sub>7</sub>G<sub>1</sub>  $\cup$  G<sub>4</sub> is a subdivision of B<sub>1</sub>G<sub>3</sub>  $\cup$  G<sub>4</sub> contains B<sub>1</sub>FIGURE 21. Kuratowski covering of  $G = \tilde{I}_{9,10}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
 $U6.6a$  and  $\tilde{I}_{9,42}^2$



$G_1 \cup G_3 \cup G_4$  contains  
 $K_8 - K_{2,3}$  and  $K_8 - K_3 \vee K_2$



$G_1 \cup G_2 \cup G_4$  contains  
 $T5.8$  and  $\tilde{I}_{9,29}^2$



$G_1 \cup G_2 \cup G_3$  contains  
 $T5.8$  and  $\tilde{I}_{9,24}^2$















FIGURE 22. Kuratowski covering of  $G = \tilde{I}_{9,10}^3$  for  $\mathbb{N}_3$  (Continued)


$$G = K_9 - \begin{array}{c} \bullet^4 \\ 5 \quad \bullet^1 \\ \downarrow \quad \downarrow \\ \bullet^6 \quad \bullet^8 \end{array}$$

G<sub>1</sub>G<sub>2</sub>G<sub>3</sub>G<sub>4</sub>G<sub>1</sub>  $\cup$  G<sub>2</sub> contains D<sub>3</sub>G<sub>2</sub>  $\cup$  G<sub>3</sub> contains F<sub>1</sub>G<sub>1</sub>  $\cup$  G<sub>3</sub> contains a subdivision of D<sub>3</sub>G<sub>2</sub>  $\cup$  G<sub>4</sub> contains D<sub>6</sub>G<sub>1</sub>  $\cup$  G<sub>4</sub> contains a subdivision of D<sub>17</sub>G<sub>3</sub>  $\cup$  G<sub>4</sub> contains D<sub>3</sub>FIGURE 23. Kuratowski covering of  $G = \tilde{I}_{9,11}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_3 \cup G_4$  contains  
S5.6 and  $\tilde{I}_{9,39}^2$



$G_1 \cup G_2 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_3$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$













FIGURE 24. Kuratowski covering of  $G = \tilde{I}_{9,11}^3$  for  $\mathbb{N}_3$  (Continued)


$$G = K_9 - \begin{array}{ccccccccc} & & & & 2 \\ & & & & | \\ & & & & 6 & 1 & 3 & 4 & 5 \end{array}$$

G<sub>1</sub>G<sub>2</sub>G<sub>3</sub>G<sub>4</sub>G<sub>1</sub> ∪ G<sub>2</sub> contains B<sub>3</sub>G<sub>2</sub> ∪ G<sub>3</sub> contains a subdivision of B<sub>1</sub>G<sub>1</sub> ∪ G<sub>3</sub> contains E<sub>20</sub>G<sub>2</sub> ∪ G<sub>4</sub> contains C<sub>7</sub>G<sub>1</sub> ∪ G<sub>4</sub> is a subdivision of B<sub>1</sub>G<sub>3</sub> ∪ G<sub>4</sub> contains a subdivision of B<sub>1</sub>FIGURE 25. Kuratowski covering of  $G = \tilde{I}_{9,12}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_4$  contains  
T5.8 and  $\tilde{I}_{9,27}^2$



$G_1 \cup G_2 \cup G_3$  contains  
T5.8 and  $\tilde{I}_{9,35}^2$






FIGURE 26. Kuratowski covering of  $G = \tilde{I}_{9,12}^3$  for  $\mathbb{N}_3$  (Continued)


$$G = K_9 - \begin{array}{c} 1 \\ \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ 3 \quad 2 \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ 5 \quad 4 \end{array}$$




G<sub>1</sub>




G<sub>2</sub>




G<sub>3</sub>




G<sub>4</sub>




$G_1 \cup G_2$  contains  $D_3$




$G_2 \cup G_3$  contains a subdivision of  $B_1$




$G_1 \cup G_3$  contains  $D_{17}$



$G_3 \cup G_4$  contains  $C_7$




$G_1 \cup G_4$  contains a subdivision of  $B_1$




$G_3 \cup G_4$  contains  $B_1$

FIGURE 27. Kuratowski covering of  $G = \tilde{I}_{9,13}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
 $V6.5$  and  $\tilde{I}_{9,32}^2$



$G_1 \cup G_3 \cup G_4$  contains  
 $K_8 - K_3$  and  $K_8 - 2K_3$



$G_1 \cup G_2 \cup G_4$  contains  
 $S5.5$  and  $\tilde{I}_{9,38}^2$



$G_1 \cup G_2 \cup G_3$  contains  
 $S5.6$  and  $\tilde{I}_{9,47}^2$










FIGURE 28. Kuratowski covering of  $G = \tilde{I}_{9,13}^3$  for  $\mathbb{N}_3$  (Continued)


$$G = K_9 -$$

G<sub>1</sub>G<sub>2</sub>G<sub>3</sub>G<sub>4</sub>


$G_1 \cup G_2$  contains a subdivision of  $B_1$        $G_2 \cup G_3$  contains a subdivision of  $B_1$



$G_1 \cup G_3$  contains  $C_7$



$G_2 \cup G_4$  contains  $C_7$



$G_1 \cup G_4$  contains a subdivision of  $B_1$        $G_3 \cup G_4$  contains a subdivision of  $B_1$






FIGURE 29. Kuratowski covering of  $G = \tilde{I}_{9,14}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
V6.4 and  $\tilde{I}_{9,41}^2$



$G_1 \cup G_3 \cup G_4$  contains  
V6.4 and  $\tilde{I}_{9,41}^2$



$G_1 \cup G_2 \cup G_4$  contains  
V6.4 and  $\tilde{I}_{9,41}^2$



$G_1 \cup G_2 \cup G_3$  contains  
V6.4 and  $\tilde{I}_{9,41}^2$







FIGURE 30. Kuratowski covering of  $G = \tilde{I}_{9,14}^3$  for  $\mathbb{N}_3$  (Continued)

FIGURE 31. Kuratowski covering of  $G = \tilde{I}_{9,15}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
 $K_8 - (K_{1,2} \cup 2K_2)$  and  $K_8 - 2K_3$



$G_1 \cup G_3 \cup G_4$  contains  
U6.6a and  $\tilde{I}_{9,39}^2$



$G_1 \cup G_2 \cup G_4$  contains V7.6 and  $\tilde{I}_{9,6}^2$



$G_1 \cup G_2 \cup G_3$  contains T5.8 and  $\tilde{I}_{9,21}^2$







FIGURE 32. Kuratowski covering of  $G = \tilde{I}_{9,15}^3$  for  $\mathbb{N}_3$  (Continued)

FIGURE 33. Kuratowski covering of  $G = \tilde{I}_{9,16}^3$  for  $\mathbb{N}_3$


$G_2 \cup G_3 \cup G_4$  contains  
 $S5.6$  and  $\tilde{I}_{9,5}^2$



$G_1 \cup G_3 \cup G_4$  contains  
 $K_8 - K_{2,3}$  and  $K_8 - (K_3 \vee K_2)$



$G_1 \cup G_2 \cup G_4$  contains  
 $U6.6a$  and  $\tilde{I}_{9,2}^2$



$G_1 \cup G_2 \cup G_3$  contains  
 $S5.6$  and  $\tilde{I}_{9,6}^2$





FIGURE 34. Kuratowski covering of  $G = \tilde{I}_{9,16}^3$  for  $\mathbb{N}_3$  (Continued)

FIGURE 35. Kuratowski covering of  $G = \tilde{I}_{9,1}^4$  for  $\mathbb{N}_4$







FIGURE 36. Kuratowski covering of  $G = \tilde{I}_{9,1}^4$  for  $N_4$  (Continued)




$G_2 \cup G_5$  contains a subdivision of  $B_1$




$G_1 \cup G_3 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$




$G_3 \cup G_4$  contains a subdivision of  $D_{17}$




$G_1 \cup G_2 \cup G_5$  contains  
 $K_8 - K_{2,3}$  and  $K_8 - 2K_{1,3}$




$G_3 \cup G_5$  contains a subdivision of  $B_1$



$G_1 \cup G_2 \cup G_4$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$





$G_4 \cup G_5$  contains  $D_3$




$G_1 \cup G_2 \cup G_3$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$

FIGURE 37. Kuratowski covering of  $G = \tilde{I}_{9,1}^4$  for  $N_4$  (Continued)


FIGURE 38. Kuratowski covering of  $G = \tilde{I}_{9,2}^4$  for  $\mathbb{N}_4$




$G_1 \cup G_4$  contains  $D_3$




$G_2 \cup G_3 \cup G_5$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$




$G_1 \cup G_5$  contains a subdivision of  $B_1$




$G_2 \cup G_3 \cup G_4$  contains  
V7.7 and  $\tilde{I}_{9,2}^2$




$G_2 \cup G_3$  contains a subdivision of  $B_1$



$G_1 \cup G_4 \cup G_5$  contains  
S5.5 and  $\tilde{I}_{9,38}^2$




$G_2 \cup G_4$  contains  $D_3$



$G_1 \cup G_3 \cup G_5$  contains  
S5.6 and  $\tilde{I}_{9,8}^2$

FIGURE 39. Kuratowski covering of  $G = \tilde{I}_{9,2}^4$  for  $N_4$  (Continued)

FIGURE 40. Kuratowski covering of  $G = \tilde{I}_{9,2}^4$  for  $N_4$  (Continued)

## REFERENCES

1. D. Archdeacon, *A Kuratowski theorem for the projective plane*, J. Graph Theory **5** (1981), 243–246.
2. D. Archdeacon and J. P. Huneke, *A Kuratowski theorem for nonorientable surfaces*, J. Combin. Theory Ser. B **46** (1989), 173–231.
3. J. Cashy, *Irreducible graphs for the Klein bottle*, Ph.D. thesis, Ohio State University, 2000.
4. R. A. Duke and G. Haggard, *The genus of subgraphs of  $K_8$* , Israel J. Math. **11** (1972), 452–455.
5. H. Glover, J. P. Huneke, and C. Wang, *103 graphs that are irreducible for the projective plane*, J. Combin. Theory Ser. B **27** (1979), 332–370.
6. A. Hlavacek, *9-vertex irreducible graphs on the torus*, Ph.D. thesis, Ohio State University, 1997.
7. J. P. Huneke, D. McQuillan, and R. B. Richter, *Orientably simple graphs of genus 1*, preprint, 1995.
8. S. Hur, *Minimal forbidden subgraphs of order  $< 10$  for the nonorientable surfaces of genus 3 and 4*, submitted for publication.
9. ———, *The Kuratowski covering conjecture for graphs of order  $< 10$* , Proceedings of the Thirty-Ninth Southeastern International Conference on Combinatorics, Graph Theory and Computing, 2008, submitted for publication.
10. ———, *The Kuratowski covering conjecture for graphs of order less than 10*, Ph.D. thesis, Ohio State University, 2008.
11. K. Kuratowski, *Sur le problème des courbes gauches en topologie*, Fund. Math. **15** (1930), 271–283.
12. B. Mohar and C. Thomassen, *Graphs on Surfaces*, Johns Hopkins University Press, 2001.
13. W. Myrvold, Personal Communication.
14. W. Myrvold and E. Neufeld, *Practical toroidality testing*, Proceedings of the Eighth Annual ACM-SIAM Symposium on Discrete Algorithms, 1997, pp. 574–580.
15. N. Robertson and P. D. Seymour, *Graph minors. VIII. A Kuratowski theorem for general surfaces*, J. Combin. Theory Ser. B **48** (1990), 255–288.

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, 231 WEST 18TH AVENUE,  
 COLUMBUS, OH 43210  
*E-mail address:* sjin@math.ohio-state.edu